

AD _____

Award Number: DAMD17-02-1-0610

TITLE: Membrane Estrogen Receptor Alpha Targeting and Its
Association with SHC in Regulating Breast Cancer Cell
Proliferation

PRINCIPAL INVESTIGATOR: Robert X-D Song, M.D., Ph.D.

CONTRACTING ORGANIZATION: University of Virginia
Charlottesville, Virginia 22904

REPORT DATE: June 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20031031 117

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 2003	3. REPORT TYPE AND DATES COVERED Annual (6 May 02 - 5 May 03)	
4. TITLE AND SUBTITLE Membrane Estrogen Receptor Alpha Targeting and Its Association with SHC in Regulating Breast Cancer Cell Proliferation			5. FUNDING NUMBERS DAMD17-02-1-0610	
6. AUTHOR(S) Robert X-D Song, M.D., Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Virginia Charlottesville, Virginia 22904 E-Mail: rs5wf@virginia.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited.				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) <p>17β-estradiol (E2) induces rapid, non-genomic effect in MCF-7 breast cancer cells, including rapid activation of MAPK, phosphorylation of adapter protein Shc, increase of the interaction between Shc and estrogen receptor (ERα). More strikingly E2 also induced a rapid membrane association of ERα (1). Further studying the structure and function of ERα, we demonstrated that only membrane-associated ERα, but not cytosol and nuclear ones, mediates E2 effect on MAPK activation (2). To study the mechanism of ERα membrane association, we further investigated the role of Shc in estrogen action. Here we demonstrated that in MCF-7 cells, Shc acts as a chaperon linking ERα to the cell membrane by binding to a transmembrane receptor IGF-1R. Further study is under the way to investigate the molecular mechanism among these protein complex formation and their biological effects in estrogen non-genomic action.</p>				
14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	5
Conclusions.....	6
References.....	6
Appendices.....	N/A

Introduction

17 β -estradiol (E2) induces rapid, non-genomic effect in MCF-7 breast cancer cells, including rapid activation of MAPK, phosphorylation of adapter protein Shc, increase of the interaction between Shc and estrogen receptor (ER α). More strikingly E2 also induced a rapid estrogen receptor membrane association (1). Further studying the structure and function of ER α , we demonstrated that only membrane-associated ER α , but not cytosol and nuclear ones, mediates E2 effect on MAPK activation (2). Since Shc physically interacts with many trans-membrane growth factor receptors, such as IGF-1 receptor (IGF-1R), EGF receptor (EGFR), PDGF receptor (PDGFR) and Insulin receptor (IR), it is conceivable that Shc might be a chaperon brings ER α to the membrane by binding to one of above receptors. So far only IGF-1R and EGFR are reported to be involved in rapid E2 action (3;4). Based on this, we reasoned that Shc might be a molecule serving two functions, one is to bring ER α to the membrane by interaction with either IGF-1R or EGFR, and the other is to activate IGF-1R-initiated MAPK signaling pathway.

Body

In Task 4 of my grant proposal, we proposed to investigate the role of adaptor protein Shc on its function to mediate ER α membrane association in MCF-7 breast cancer cells. Shc is a common

substrate of many transmembrane growth factor receptors, such as IGF-1R, IR, EGFR and PDGFR, activation of these receptor will lead to Shc physically association with these membrane receptors. So far only IGF-1R and EGFR are reported to be involved in rapid E2 action (3;4). Accordingly, we tested several protein expression, such as IGF-1R, EGFR, ER α and Shc, in MCF-7, LTED and COS-1 cells. LTED cells were developed from MCF-7 cells by long-term estrogen-deprivation and characterized as supersensitive to E2 treatment than their parental cells due to elevated ER α levels (5;6). Interestingly, all cells express IGF-1R from medium to high levels and only COS-1 cells, but not MCF-7 and LTED cells, show the EGFR expression, indicating that IGF-1R, but not




Fig. 1. Expression of proteins in MCF-7, LTED and COS-1 cells.

ERGR, might be a potential transmembrane receptor interacting with Shc (Fig. 1). To demonstrate that Shc plays an important role in mediating ER α membrane association, a gene silencing method was employed by using siRNA against Shc. We transfected a pool of 4 siRNA's, each which were designed to knock down Shc. We also used a




Fig. 2. Down-regulation of Shc using siRNA. MCF-7 cells were grown for 2 days after transfection of the siRNA against Shc. A Western blot was performed using cell lysates. The PVDF membrane was blotted with anti-Shc, anti-ER α , anti-IGF1R, anti-IRS1 and anti-VDR antibodies (left panels). The specificity of siRNA against Shc was also tested using confocal microscopy. Cells transfected with or without Shc were immunofluorescently stained with anti-Shc (blue) and anti-ER α (green). The images were taken under the same scale.

control pool of irrelevant siRNA molecules as negative control. Both siRNA were purchased from Dharmacon Inc. As shown in Fig. 2 (left panels), expression of siRNA against Shc exclusively knocked down the Shc protein without altering the levels of any other proteins (ER α , IGF-1R, IRS-1 and VDR) tested in this experiment. The specificity of Shc siRNA was also further confirmed by confocal microscopy method, showing that down-regulation of Shc did not change ER α expression in the cells (right panels in Fig. 2). The interaction between ER α and IGF-1R has been reported in ER α -transfected COS-1 cells (3). Since we demonstrated that ER α physically interacted with Shc, we were wondering if both ER α and IGF-1R interaction is Shc-dependent.

To do so, one of our strategies is to knock down Shc in MCF-7 cells and test if this can alter the IGF-1R and ER α association.

Fig. 3. shows that siRNA against Shc decreased Shc protein expression in MCF-7 cells (lower panel).

Estrogen treatment increased the ER α and IGF-1R interaction in control siRNA expressed cells. Down-regulation of Shc blocked ER α interaction with IGF-1R,

indicating that Shc is an intermediate protein linking both ER α and IGF-1R association. Further study on the functions of this triple protein formation is currently under active investigation.




Fig. 3. Knocking down Shc blocked the interaction between ER α and IGF-1R. Cells were transfected with or without siRNA against Shc. The protein levels of Shc was shown in Western blot (lower panel). The ER α and IGF-1R interaction was measured by immunoprecipitation of IGF-1R and detection of ER α on Western blot (upper panel).

Key research accomplishments

1. We successfully constructed an ER α -expression vector by ligation of DNA sequence coding for a membrane-signaling sequence of GAP-43 protein on the C-terminal of ER α insert (Task 1) (2).
2. Functional testing the expressed membrane-targeted ER α has been done using confocal immunofluorescence microscopy. Comparing with wild type ER α that show 99% expressed in the nucleus, our data show that expression this vector in COS-1 cells lead to 45% of ER α expressed on the cell membrane and rest of it in the cytosol (Task 2) (2).
3. We then tested the ER α -mediated MAPK activation in COS-1 cells. Compared with the wild type one (99% in nucleus), Cytosol (99% in cytosol), the membrane expressed ER α is the only form that mediated estrogen effect on MAPK phosphorylation (Task 3) (2).
4. We are currently working on the mechanism how adapter protein Shc mediates ER α on its membrane association and MAPK activation.

Reportable outcomes:

- 1) We have one paper published under this grant supporting mechanism. Title: Membrane association of estrogen receptor α mediates estrogen effect on MAPK activation. BBRC, 294: 926-933, 2002 (2).

- 2) A manuscript is under writing, entitled "Adapter protein Shc as a chaperon mediates estrogen receptor membrane association in breast cancer cells".
- 3) Three abstracts were submitted to Endo2003 in Philadelphia, PA
 - 1, IGF-1 receptor activation and its physical interactions with adapter protein Shc and ER alpha mediate the non-genomic action of estradiol in breast cancer cells. Song, RX, Zhang, Z, Boa, Y, Black, MJ, and Santen, RJ. Department of internal medicine, UVA, Charlottesville, VA 22903
 2. Antiestrogen and estrogens regulate cell proliferation and apoptosis differently in long-term estrogen-deprived and wild type MCF-7 cells. Zhang, Z, Boa, Y, Black, MJ, Santen, RJ and Song, RX, Department of Internal Medicine, UVA, Charlottesville, VA 22903
 3. MAP kinase negatively regulates estrogen receptor-mediated transcriptional activity in human breast cancer cells. Zhang, Z, Boa, Y, Black, MJ, Santen, RJ and Song, RX, Department of Internal Medicine, UVA, Charlottesville, VA 22903
- 4) The results from this grant was presented in the following meetings:
 1. Invited speaker in one of symposiums "Recognition of estrogen receptor in estrogen non-genomic action". Title: Involvement of Shc in estrogen receptor membrane association in breast cancer cells. FASEB – Cell Biology, San Diego, 4-14-2003.
 2. Invited speaker. Title: Upstream signaling pathways in estrogen action on MAPK activation in breast cancer cells. Department of Biochemistry, University of California – Riverside, 4-16-2003.
 3. Invited speaker. Title: Upstream signaling pathways in estrogen action on MAPK activation in breast cancer cells. Department of Microbiology of UVA, Charlottesville, VA, 4-8-2003.
 4. Presenting in Division of Endocrinology, Department of internal medicine, UVA, Charlottesville, 6-3-2003. Title: Non-genomic signaling pathway of estrogen and breast cancer.

Conclusions

Adapter protein Shc acts as a chaperon and it involves in rapid estrogen action linking ER α to the cell membrane by binding on IGF-1 receptor.

Reference List

1. Song RX, McPherson RA, Adam L, Bao Y, Shupnik M, Kumar R, Santen RJ 2002 Linkage of Rapid Estrogen Action to MAPK Activation by ER α -Shc Association and Shc Pathway Activation. *Mol Endocrinol* 16:116-127

2. Zhang Z, Maier B, Santen RJ, Song R.X 2002 Membrane association of estrogen receptor alpha mediates estrogen effect on MAPK activation. *Biochem Biophys Res Commun* 294:000
3. Kahlert S, Nuedling S, van Eickels M, Vetter H, Meyer R, Grohe C 2000 Estrogen receptor alpha rapidly activates the IGF-1 receptor pathway. *J Biol Chem* 275:18447-18453
4. Richards RG, DiAugustine RP, Petrusz P, Clark GC, Sebastian J 1996 Estradiol stimulates tyrosine phosphorylation of the insulin-like growth factor-1 receptor and insulin receptor substrate-1 in the uterus. *Proc Natl Acad Sci U S A* 93:12002-12007
5. Martin LA, Farmer I, Johnston SR, Ali S, Marshall CJ, Dowsett M 2003 Enhanced ERalpha ERBB2 and MAPK signal transduction pathways operate during the adaptation of MCF-7 cells to long term oestrogen deprivation. *J Biol Chem*
6. Santen R, Jeng MH, Wang JP, Song R, Masamura S, McPherson R, Santner S, Yue W, Shim WS 2001 Adaptive hypersensitivity to estradiol: potential mechanism for secondary hormonal responses in breast cancer patients. *The Journal of Steroid Biochemistry and Molecular Biology* 79:115-125